


WHAT IS CLAIMED IS:

1. A cosmetic method of skin lightening comprising applying to the skin a composition comprising:
- 5 a. about 0.000001 to about 50 % of a compound of general formula I (I)




- Where each or both R₁ and/or R₂ represents hydrogen (H); linear or branched, saturated or unsaturated C₁ – C₁₂ alkyl, alkenyl, acyl, or heteroalkyl groups;
- 10 R₃ represents linear or branched, cyclic or acyclic, saturated or unsaturated C₁ – C₁₂ alkyl, alkenyl, cycloalkyl, cycloalkenyl, or heteroalkyl group;
- R₄ represents a hydrogen atom (H); straight or branched, cyclic or acyclic, saturated or unsaturated, containing or not containing a heteroatom C₁-C₂₂ alkyl, alkenyl, cycloalkyl, cycloalkenyl, heteroalkyl, aryl, or heteroaryl group;
- 15 and
- b. a cosmetically acceptable carrier.

- 20 2. The method of claim 1, wherein said composition further comprises a sunscreen.

3. The method of claim 2, wherein said sunscreen is a micronized metal oxide.
4. The method of claim 1, wherein said compound is a 4-methyl 7-hydroxy coumarin derived resorcinol derivative.
5. The method of claim 1, wherein said compound is a compound of formula II:


(II)



6. The cosmetic method of claim 1, wherein R_1 and R_2 both represent hydrogen.
7. The cosmetic method according to claim 1, wherein said composition further comprises a skin benefit agent selected from the group consisting of alpha-hydroxy acids, beta-hydroxy acids, polyhydroxy acids, hydroquinone, t-butyl hydroquinone, Vitamin C derivatives, dioic acids, retinoids, resorcinol derivatives, and mixtures thereof.
8. The cosmetic method of claim 1, wherein said composition further comprises an organic sunscreen selected from the group consisting of Benzophenone-3, Benzophenone-4, Benzophenone-8, DEA, Methoxycinnamate, Ethyl

dihydroxypropyl-PABA, Glyceryl PABA, Homosalate, Methyl anthranilate, Octocrylene, Octyl dimethyl PABA, Octyl methoxycinnamate (PARSOL MCX), Octyl salicylate, PABA, 2-Phenylbenzimidazole-5-sulphonic acid, TEA salicylate, 3-(4-methylbenzylidene)-camphor, Benzophenone-1,
5 Benzophenone-2, Benzophenone-6, Benzophenone-12, 4-Isopropyl dibenzoyl methane, Butyl methoxy dibenzoyl methane (PARSOL 1789), Octocrylene, and mixtures thereof.

9. A cosmetic composition comprising:
10 a. about 0.000001 to about 50 % of a compound of general formula I:
(I)




- wherein each or both R₁ and/or R₂ represents hydrogen (H); linear or branched, saturated or unsaturated C₁ – C₁₂ alkyl, alkenyl, acyl, or heteroalkyl groups;
15 R₃ represents linear or branched, cyclic or acyclic, saturated or unsaturated C₁ – C₁₂ alkyl, alkenyl, cycloalkyl, cycloalkenyl, or heteroalkyl group;
R₄ represents a hydrogen atom (H); straight or branched, cyclic or acyclic,
20 saturated or unsaturated, containing or not containing a heteroatom C₁-C₂₂ alkyl, alkenyl, cycloalkyl, cycloalkenyl, heteroalkyl, aryl, or heteroaryl group; and
b. a cosmetically acceptable carrier.

10. The cosmetic composition of claim 9, wherein said compound is a compound of formula II:

(II)

5



11. The cosmetic composition of claim 9, wherein R₁ and R₂ both represent hydrogen.

10


12. The cosmetic composition of claim 9, wherein said compound comprises about 0.00001 % to about 10 % of said composition.

13. The cosmetic composition of claim 9, wherein said compound comprises about 0.001 % to about 7 % of said composition.

15

14. The cosmetic composition of claim 9, wherein said compound comprises about 0.01 % to about 5 % of said composition.

15. A process for making compounds having a general formula selected from the group consisting of B, C, D, and mixtures thereof, comprising:



5


wherein

R_3 represents linear or branched, cyclic or acyclic, saturated or unsaturated $C_1 - C_{12}$ alkyl, alkenyl, cycloalkyl, cycloalkenyl, or heteroalkyl group;

10 R_4 represents a hydrogen atom (H); straight or branched, cyclic or acyclic, saturated or unsaturated, containing or not containing a heteroatom $C_1 - C_{22}$ alkyl, alkenyl, cycloalkyl, cycloalkenyl, heteroalkyl, aryl, or heteroaryl group.

16. The process of claim 15 further comprising substitution of the 1,3-hydroxy positions of the phenyl ring to yield compound of general formula I:


(I)



5 wherein each or both R₁ and/or R₂ represents hydrogen (H); linear or branched, saturated or unsaturated C₁ – C₁₂ alkyl, alkenyl, acyl, or heteroalkyl groups.

17. A compound of general formula I

(I)



10

Where each or both R₁ and/or R₂ represents hydrogen (H); linear or branched, saturated or unsaturated C₁ – C₁₂ alkyl, alkenyl, acyl, or heteroalkyl groups;

R₃ represents linear or branched, cyclic or acyclic, saturated or unsaturated C₁ –
15 C₁₂ alkyl, alkenyl, cycloalkyl, cycloalkenyl, or heteroalkyl group;


R₄ represents a hydrogen atom (H); straight or branched, cyclic or acyclic, saturated or unsaturated, containing or not containing a heteroatom C₁-C₂₂ alkyl, alkenyl, cycloalkyl, cycloalkenyl, heteroalkyl, aryl, or heteroaryl group.

5

18. The compound of claim 17, wherein said compound is a compound of formula II:

(II)

10



15 19. The compound of claim 1, wherein said compound is a 4-methyl 7-hydroxy coumarin derived resorcinol derivative.

20. The compound of claim 1, wherein R₁ and R₂ both represent hydrogen.